AD-A276 232

3 August 1993

GE

Form Approved OBM No. 0704-0188

:luding the time for reviewing instructions, searching existing data sources, gathering and garding this burden or any other aspect of this collection of information, including suggestions nd Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to

3. Report Type and Dates Covered. Contractor Report

4. Title and Subtitle.

6. Author(s).

Public reporting burden for

maintaining the data needed

for reducing this burden, to

the Office of Management a

Penetrable Wedge Analysis

1. Agency Use Only (Luaro biatin).

5. Funding Numbers.

Contract

N00014-93-1-6014

Program Element No. 0601153N

Project No.

Task No.

Accession No.

DN257115


Work Unit No.

12212F

7. Performing Organization Name(s) and Address(es).

Robert W. Scarstein* and Anthony M. J. Davis*

*The University of Alabama Tuscaloosa, AL 35487-0286



8. Performing Organization Report Number.

9. Sponsoring/Monitoring Agency Name(s) and Address(es).

Naval Research Laboratory Acoustic Simulation and Tactics Branch Stennis Space Center, MS 39529-5004

10. Sponsoring/Monitoring Agency Report Number.

NRL/CR/7181--93-0001

11. Supplementary Notes.

12a. Distribution/Availability Statement.

Approved for public release; distribution is unlimited.

12b. Distribution Code.

13. Abstract (Maximum 200 words).

The continuation of this wedge scattering research is proceeding as proposed, with several significant accomplishments and findings to date. The nature of this rigorous mathematical approach to scattering problems is such that original formulation is very important and usually very slow to evolve. For example, the previous work summarized in the report [61] had several abandoned and potentially useful starts, while the finished product is substantial.

94-03651 1 P95

14. Subject Terms.

Scattering, plane waves

15. Number of Pages. 6

16. Price Code.

17. Security Classification of Report.

Unclassified

18. Security Classification of This Page. Unclassified

19. Security Classification of Abstract. Unclassified

20. Limitation of Abstract.

SAR

Approved for public release; distribution is unlimited.

Quarterly Progress Report Submitted to:
US Naval Research Laboratory - Stennis Space Center

PENETRABLE WEDGE ANALYSIS GRANT NO. N00014-93-1-6014

ROBERT W. SCHARSTEIN, PRINCIPAL INVESTIGATOR DEPARTMENT OF ELECTRICAL ENGINEERING

Accesion for

NTIS CRA&I
DITIC TAB

Unit of the distribution of th

AND

ANTHONY M. J. DAVIS, Co-PRINCIPAL INVESTIGATOR DEPARTMENT OF MATHEMATICS

The University of Alabama Tuscaloosa, Alabama 35487-0286 205-348-1761

DTIC QUALITY INSPECTED &

3 August 1993

The continuation of this wedge scattering research is proceeding as proposed, with several significant accomplishments and findings to date. The nature of this rigorous mathematical approach to scattering problems is such that original formulation is very important and usually very slow to evolve. For example, the previous work summarized in the report [61] had several abandoned and potentially useful starts, while the finished product is substantial.

I. JOURNAL PAPER

A journal paper titled "Mellin Transform Solution for the Static Line-Source Excitation of a Dielectric Wedge" has been submitted, revised, and accepted for publication in the IEEE Transactions on Antennas and Propagation [62]. A conference paper also summarizing these results was presented in July 1993 at the 1993 Radio Science Meeting in Ann Arbor, MI.

II. GENERAL PENETRABLE WEDGE

In order to make the most significant progress possible, our initial philosophy during this first quarter of this research has been to concentrate on the most difficult aspect - the truly arbitrary, penetrable wedge. The solution to the impedance boundary wedge could then be extracted from the more general (and valuable) result. I have had scientific dialog with two leading scattering theorists, Egon Marx [20-22,28] and Ismo Lindell [63], who

have made recent contributions to this and related wedge scattering problems. I would summarize the current (August 1993) thinking on the problem to be:

- (1) The penetrable wedge problem remains one of the great unsolved canonical scattering geometries.
- (2) Numerical "solutions" are still proliferating, but are too messy and removed from the physics to offer any important insight into the wave mechanisms that we hope to uncover and understand.

Based on last year's work [61] with the Kontorovich-Lebedev transform, which is a superposition of cylindrical waves, a general superposition of arbitrary plane waves has been investigated as a suitable field representation. Such a (spatial) spectrum of propagating $(|k_x| < k_j)$ and evanescent $(|k_x| > k_j)$ plane waves

$$\psi_j(x,y) = \int_{-\infty}^{\infty} dk_x \, \Psi_j(k_x,k_y) e^{i(k_x x + k_y y)}$$

where $k_x^2 + k_y^2 = k_j^2$ and j = 1, 2 denotes the two different media, may work in principle. A rather novel approach is to apply a conformal transformation to the homogeneous wedge region to obtain an effectively inhomogeneous half-space region. This results in a spatially varying wavenumber appearing in the transformed Helmholtz equation

$$(\nabla^2 + \kappa^2(\vec{r})) \psi(\vec{r}) = 0.$$

This approach demonstrates all too well a conservation of work principle: the problem can be cast in terms of the nice (familiar) Helmholtz operator with troublesome wedge boundaries or as a simple planar boundary that constrains a messy partial differential operator.

The lack of a rigorous solution technique to apply to the truly arbitrary wedge is disappointing to us, and is probably a posteriori expected by some colleagues. Fortunately, we can now confidently proceed with a new appreciation and respect for our primary goal of this year's research project – the mathematical solution of the impedance boundary wedge.

III. PHYSICAL IMPEDANCE BOUNDARY CONDITION

The coupled difference equations (14), (17), and (18) on page 27 of the previous report [61] are solved by first eliminating the function $C(\nu)$:

$$A(\nu)\sin(\nu\pi) + B(\nu)\cos(\nu\pi) = \cos[\nu(\phi'-\pi)]$$

$$\frac{k\eta}{2} \left\{ A(\nu - 1)\sin[(\nu - 1)\alpha] + B(\nu - 1)\cos[(\nu - 1)\alpha] + A(\nu + 1)\sin[(\nu + 1)\alpha] + B(\nu + 1)\cos[(\nu + 1)\alpha] \right\} - A(\nu)\cos(\nu\alpha) + B(\nu)\sin(\nu\alpha) = 0.$$

Application of an integral operator, such as the Laplace or Mellin inversion formula, eliminates the troublesome $\nu-1$ and $\nu+1$ and yields purely algebraic equations in the coefficient functions $A(\nu)$ and $B(\nu)$. Finding such a transformation that also results in explicit analytic expressions that can be conveniently carried through the inverse Kontorovich-Lebedev transform is an arduous task. Our present status is that of completed inversions via both the Laplace and the Mellin formulae, with results to date partially encouraging but still too complicated. We have identified several alternatives, including an asymptotic approach that lets $\nu=i\tau$ as it appears in the Kontorovich-Lebedev inversion, and concentrating on large τ . Adding to our motivation is the comforting knowledge that our well-defined boundary value problem possesses a unique solution which is defined in a single, homogeneous region of space. Also, the formulation and execution of the mathematics is rigorous and complete.

IV. INHOMOGENEOUS OR PSEUDO-IMPEDANCE BOUNDARY

The source of the shifting of ν to $\nu-1$ and $\nu+1$ in the above case of the Leontovich boundary condition

$$\psi(r,\alpha) + \frac{\eta}{r} \frac{\partial}{\partial \phi} \psi(r,\alpha) = 0$$

is the 1/r factor from the normal derivative

$$\frac{\partial}{\partial n} = \frac{1}{r} \frac{\partial}{\partial \phi}.$$

The observation on page 28 of [61] that an inhomogeneous surface impedance that is proportional to radius

$$\eta = r \eta'$$

avoids any difference equation in the transform coefficient functions. In this special case, much asymptotic progress and physical interpretation can be accomplished, as indicated in [61]. I did not emphasize this particular avenue, since it appears to be a problem contrived to fit a solution, rather than the solution to a real physical problem. Although Felsen and Marcuvitz [65] clearly recognized the analytic attractiveness of this "pseudo-impedance" wedge circa 1973, I only recently (today!) discovered their passing mention of it. They make no apology at all for any detachment from "reality." Perhaps I should not be so quick to dismiss this curious variation on the impedance boundary wedge. In any event, we can always return to this easier problem at a later date.

REFERENCES

- 1. A.D. Rawlins, Diffraction by a Dielectric Wedge, J. Inst. Maths Applies 19 (1977), 261-279.
- 2. S. Berntsen, Diffraction of an Electric Polarized Wave by a Dielectric Wedge, SIAM J. Appl. Math. 43 (1983), 186-211.
- 3. V. Rokhlin, Solution of Acoustic Scattering Problems by Means of Second Kind Integral Equations, Wave Motion 5 (1983), 257-272.
- 4. R.E. Kleinman and P.A. Martin, On Single Integral Equations for the Transmission Problem of Acoustics, SIAM J. Appl. Math. 48 (1988), 307-325.
- 5. K. Davey, An Integral Approach to Electromagnetic Scattering off a Dielectric Wedge, Electromagnetics 7 (1987), 167-183.
- 6. G.D. Maliuzhinets, Excitation, Reflection and Emission of Surface Waves from a Wedge with Given Face Impedances, Sov. Phys. Dokl. 3 (1958), 752-755.
- W.E. Williams, Diffraction of an E-polarized Plane Wave by an Imperfectly Conducting Wedge, Proc. R. Soc. London Ser. A 252 (1959), 376-393.
- 8. T.B.A. Senior, Diffraction by an Imperfectly Conducting Wedge, Commun. Pure Appl. Math. 12 (1959), 337-372.
- 9. A.D. Pierce and W.J. Hadden, Jr., Plane wave diffraction by a wedge with finite impedance, J. Acoust. Soc. Am. 63 (1978), 17-27.
- 10. R. Tiberio, G. Pelosi, and G. Manara, A Uniform GTD Formulation for the Diffraction by a Wedge with Impedance Faces, IEEE Trans. Antennas Propagat. 33 (1985), 867-873.
- 11. K. Hongo and E. Nakajima, Polynomial Approximation of Maliuzhinets' Function, IEEE Trans. Antennas Propagat. 34 (1986), 942-947.
- 12. M.I. Herman, J.L. Volakis, and T.B.A. Senior, Analytic Expressions for a Function Occurring in Diffraction Theory, IEEE Trans. Antennas Propagat. 35 (1987), 1083-1086.
- 13. T. Griesser and C.A. Balanis, Reflections, Diffractions, and Surface Waves for an Interior Impedance Wedge of Arbitrary Angle, IEEE Trans. Antennas Propagat. 37 (1989), 927-935.
- 14. H. Uberall, Radar Scattering from Imperfectly Conducting or Coated Perfectly Conducting Wedges and Cones, Acta Physica Austriaca 22 (1966), 67-93.
- 15. T.S. Angell and R.E. Kleinman, Scattering of Acoustic Waves by Impedance Surfaces, Wave Phenomena: Modern Theory and Applications (C. Rogers and T.B. Moodie, eds.), Elsevier, Amsterdam, 1984, pp. 329-336.
- 16. J. Bach Andersen and V.V. Solodukhov, Field Behavior Near a Dielectric Wedge, IEEE Trans. Antennas Propagat. 26 (1978), 598-602.
- 17. S.Y. Kim, J.W. Ra, and S.Y. Shin, Edge Diffraction by Dielectric Wedge of Arbitrary Angle, Electronics Letters 19 (1983), 851-853.
- 18. P.V. Tret'yakov, Integral Solutions of the Wave Equation and the Diffraction of an Arbitrary Acoustic Wave by a Wedge, J. Appl. Maths Mechs 55 (1991), 201-205.
- 19. A.W. Glisson, An Integral Equation for Electromagnetic Scattering from Homogeneous Dielectric Bodies, IEEE Trans. Antennas Propagat. 32 (1984), 173-175.
- 20. E. Marx, Integral Equation for Scattering by a Dielectric, IEEE Trans. Antennas Propagat. 32 (1984), 166-172.
- 21. E. Marx, Computed Fields Near the Edge of a Dielectric Wedge, IEEE Trans. Antennas Propagat. 38 (1990), 1438-1442.
- 22. E. Marx, Electromagnetic Scattering from a Dielectric Wedge and the Hypersingular Integral Equation, IEEE Trans. Antennas Propagat. (1992) (submitted).
- 23. S.Y. Kim, J.W. Ra, and S.Y. Shin, Diffraction by an Arbitrary-Angled Dielectric Wedge: Part I-Physical Optics Approximation, IEEE Trans. Antennas Propagat. 39 (1991), 1272-1281.
- 24. _____, Part II-Correction to Physical Optics Solution, IEEE Trans. Antennas Propagat. 39 (1991), 1282-1292.
- 25. H.H. Syed and J.L. Volakis, Equivalent Current Formulation for an Impedance Wedge of Arbitrary Included Angle, IEEE-APS Symposium Digest (Chicago) (1992), 1857-1860.
- 26. H.H. Syed and J.L. Volakis, An Approximate Skew Incidence Diffraction Coefficient for an Impedance Wedge, Electromagnetics 12 (1992), 33-55.

- 27. C. Bergljung and L.G. Olsson, A Comparison of Solutions to the Problem of Diffraction of a Plane Wave by a Dielectric Wedge, IEEE-APS Symposium Digest (Chicago) (1992), 1861-1864.
- 28. E. Marx, The Hypersingular Integral Equation and the Dielectric Wedge, IEEE-APS Symposium Digest (Chicago) (1992), 1865-1868.
- 29. L.B. Felsen, Diffraction of the Pulsed Field from an Arbitrarily Oriented Electric or Magnetic Dipole by a Perfectly Conducting Wedge, SIAM J. Appl. Math. 26 (1974), 306-312.
- 30. J.B. Keller and A. Blank, Diffraction and Reflection of Pulses by Wedges and Corners, Commun. Pure Appl. Math. IV (1951), 75-94.
- 31. T.W. Veruttipong, Time Domain Version of the Uniform GTD, IEEE Trans. Antennas Propagat. 38 (1990), 1757-1764.
- 32. J.M. Arnold and L.B. Felsen, Rays and local modes in a wedge-shaped ocean, J. Acoust. Soc. Am. 78 (1983), 1105-1119.
- 33. A. Kamel and L.B. Felsen, Spectral theory of sound propagation in an ocean channel with weakly sloping bottom, J. Acoust. Soc. Am. 73 (1983), 1120-1130.
- 31. J.M. Arnold and L.B. Felsen, Intrinsic modes in a nonseparable ocean waveguide, J. Acoust. Soc. Am. 76 (1984), 850-860.
- 35. J.M. Arnold and L.B. Felsen, Coupled mode theory of intrinsic modes in a wedge, J. Acoust. Soc. Am. 79 (1986), 31-40.
- 36. A.P. Ansbro and J.M. Arnold, Calculation of the Green's function for the wedge-shaped layer, J. Acoust. Soc. Am. 90 (1991), 1539-1546.
- 37. J.M. Dunn, Lateral Wave Propagation in a Wedge Shaped Region, IEEE Trans. Antennas Propagat. 35 (1987), 947-955.
- 38. M.A. Biot and I. Tolstoy, Formulation of wave propagation in infinite media by normal coordinates with an application to diffraction, J. Acoust. Soc. Am. 29 (1957), 381-391.
- 39. I. Tolstoy, Diffraction by a Hard Truncated Wedge and a Strip, IEEE Trans. Oceanic Eng. 14 (1989), 4-16.
- 40. I. Tolstoy, Exact, explicit solutions for diffraction by hard sound barriers and seamounts. J. Acoust. Soc. Am. 85 (1989), 661-669.
- 41. R.S. Keiffer and J.C. Novarini, A Wedge Assemblage Method for 3-D Acoustic Scattering from Sea Surfaces: Comparison with a Helmholtz-Kirchhoff Method, Computational Acoustics Volume 1 (D. Lee, A Cakmak, and R. Vichnevetsky, eds.), Elsevier, Amsterdam, 1990, pp. 67-81.
- 42. H. Medwin, Shadowing by finite noise barriers, J. Acoust. Soc. Am. 69 (1981), 1060-1064.
- 43. A.I. Papadopoulos and C.G. Don, A study of barrier attenuation by using acoustic pulses, J. Acoust. Soc. Am. 90 (1991), 1011-1018.
- 44. C.G. Don, Application of a hard truncated wedge theory of diffraction to wide barriers, J. Acoust. Soc. Am. 90 (1991), 1005-1010.
- 45. D. Chu, Impulse response of density contrast wedge using normal coordinates, J. Acoust. Soc. Am. 86 (1989), 1883-1896.
- 46. G.B. Deane and C.T. Tindle, A three-dimensional analysis of acoustic propagation in a penetrable wedge slice, J. Acoust. Soc. Am. 92 (1992), 1583-1592.
- 47. E.K. Westwood, Broadband modeling of the three-dimensional penetrable wedge, J. Acoust. Soc. Am. 92 (1992), 2212-2222.
- 48. D.S. Jones, Methods in Electromagnetic Wave Propagation, p. 718, Clarendon Press, Oxford, 1979.
- 49. D.S. Jones, Acoustic and Electromagnetic Waves, pp. 585-590, Clarendon Press, Oxford. 1986.
- 50. W.R. Smythe, Static and Dynamic Electricity, pp. 70-72, McGraw Hill, New York, 1968.
- 51. K. Atkinson, The Numerical Solution of Laplace's Equation on a Wedge, IMA J. Num. Anal. 4 (1984), 19-41.
- 52. L.C. Andrews and B.K. Shivamoggi, Integral Transforms for Engineers and Applied Mathematicians, pp. 245-273, Macmillan. New York, 1988.
- 53. N.N. Lebedev, I.P. Skalskaya, and Y.S. Uflyand, Worked Problems in Applied Mathematics, p. 192, Dover, New York, 1965.
- 54. B. Davies, Integral Transforms and Their Applications, pp. 195-213. Springer-Verlag. New York, 1985.

- 55. J. Van Bladel, Singular Electromagnetic Fields and Sources, pp. 149-154, Clarendon Press, Oxford, 1991.
- 56. User's Manual, IMSL Math/Library: Fortran Subroutines for Mathematical Applications, pp. 767-769, IMSL Customer Relations, Houston, TX, 1989.
- 57. A.M.J. Davis and R.W. Scharstein, Electromagnetic Plane Wave Excitation of an Open-Ended, Finite-Length Conducting Cylinder, J. Electromagn. Waves Applications 7 (1993), 301-319.
- 58. R.W. Scharstein, Acoustic scattering by an open ended hard circular tube of finite length, J. Acoust. Soc. Am. 92 (1992), 3337-3342.
- 59. D. Colton and R. Kress, Integral Equation Methods in Scattering Theor., Wiley, New York, 1983.
- 60. Se-Yun Kim, Diffraction Coefficients and Field Patterns of Obtuse Angle Dielectric Wedge Illuminated by E-Polarized Plane Wave, IEEE Trans. Antennas Propagat. 40 (1992), 1427-1431.
- 61. R.W. Scharsteir. Acoustic or Electromagnetic Scattering from the Penetrable Wedge, BER Report No. 583-162, Naval Research Laboratory Stennis Space Center Contract No. N00014-92-C-6004, The University of Alabama, Tuscaloosa, AL, 1993.
- 62. R.W. Scharstein, Mellin Transform Solution for the Static Line-Source Excitation of a Dielectric Wedge, IEEE Trans. Antennas Propagat. (to appear).
- 63. M.E. Ermutlu, I.V. Lindell, and K.I. Nikoskinen, Two-dimensional Image Theory for the Conducting Wedge, Technical Report (1992), Helsinki University of Technology, Electromagnetics Laboratory, Espoo, Finland.
- 64. G.B. Deane and M.J. Buckingham, An analysis of the three-dimensional sound field in a penetrable wedge with a stratified fluid or elastic basement, J. Acoust. Soc. Am. 93 (1993), 1319-1328.
- 65. L.B. Felsen and N. Marcuvitz, Radiation and Scattering of Waves, pp. 674-675, Prentice-Hall Englewood Cliffs, NJ, 1973.